



BRIEF REPORT

## Bioprospecting of antimicrobial activity of extracts of endophytic fungi from *Bauhinia guianensis*



Eduardo A.A. Pinheiro, Jeferson R.S. Pina, André O. Feitosa, Josiwander M. Carvalho, Fábio C. Borges, Patrícia S.B. Marinho, Andrey M.R. Marinho\*

Universidade Federal do Pará, Instituto de Ciências Exatas e Naturais – Programa de Pós-graduação em Química, Rua Augusto Corrêa, 01 – Guamá, 66075-110 Belém, Pará, Brazil

Received 10 January 2016; accepted 19 August 2016  
Available online 13 January 2017

### KEYWORDS

Monocerin;  
Antimicrobial activity;  
Endophytic fungi;  
*Bauhinia guianensis*

**Abstract** Antibiotic resistance results in higher medical costs, prolonged hospital stays and increased mortality and is rising to dangerously high levels in all parts of the world. Therefore, this study aims to search for new antimicrobial agents through bioprospecting of extracts of endophytic fungi from *Bauhinia guianensis*, a typical Amazonian plant used in combating infections. Seventeen (17) fungi were isolated and as result the methanolic extract of the fungus *Exserohilum rostratum* showed good activity against the bacteria tested. The polyketide monocerin was isolated by the chromatographic technique, identified by NMR and MS, showing broad antimicrobial spectrum.

© 2016 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### PALABRAS CLAVE

Monocerin;  
Actividad antimicrobiana;  
Hongos endófitos;  
*Bauhinia guianensis*

**Bioprospección de la actividad antimicrobiana de los extractos de los hongos endófitos de *Bauhinia guianensis***

**Resumen** La resistencia a los antibióticos conduce a mayores costos médicos, hospitalizaciones prolongadas e incremento de la mortalidad, y está aumentando a niveles peligrosamente altos en todas partes del mundo. Este estudio tuvo como objetivo la búsqueda de nuevos agentes antimicrobianos a través de la bioprospección de extractos de hongos endófitos de *Bauhinia guianensis*, una planta amazónica típica, utilizada en la lucha contra problemas infecciosos. Fueron aislados 17 hongos; el extracto metanólico del hongo *Exserohilum rostratum* mostró buena actividad contra las bacterias probadas. Se aisló monocerina policétido por la técnica

\* Corresponding author.

E-mail address: [andrey@ufpa.br](mailto:andrey@ufpa.br) (A.M.R. Marinho).

<http://dx.doi.org/10.1016/j.ram.2016.08.005>

0325-7541/© 2016 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

de cromatografía; este compuesto fue identificado por RM y EM, y mostró un amplio espectro antimicrobiano.

© 2016 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Antibiotics are medicines used to prevent and treat bacterial infections. Antibiotic resistance occurs when bacteria change in response to the use of these medicines, resulting in higher medical costs, prolonged hospital stays and increased mortality and rising to dangerously high levels in all parts of the world. New resistance mechanisms emerge and spread globally every day, threatening our ability to treat common infectious diseases. A growing list of infections such as pneumonia, tuberculosis, blood poisoning and gonorrhoea are becoming harder and sometimes impossible to treat as antibiotics become less effective<sup>14</sup>.

Currently there is growing interest in endophytic microorganisms, which, in association with their host, are known to produce a wide range of compounds with diverse biological activities<sup>3,4</sup>. There are several reports in the literature of compounds isolated from endophytic fungi having antimicrobial activity, such as the extract of fungus *Nodulisporium* sp. (Xylariaceae), isolated from the woody plant species *Erica arborea* (Ericaceae), from which nodule purine compounds D, E and F that act as antifungal, antibacterial and algicide were isolated<sup>7</sup>. Of the endophytic fungus *Ampelomyces* sp. isolated from the medicinal plant *Urospermum picroides* (Asteraceae) compounds 6-O-metilalaternina and alter-solanol A were isolated, both having antimicrobial activity against gram-positive pathogens *Staphylococcus aureus*, *Staphylococcus epidermidis* and *Enterococcus faecalis*<sup>2</sup>.

In the Brazilian Amazon it is common to use prepared plants to combat various diseases. *Bauhinia guianensis* is a typical Amazon plant popularly known as "ladder tortoise", "liana toenail of the ox", "nail of the cow" and "mororó liana", which is used as a drug in folk medicine, specifically to fight infections, painful processes and diabetes. The phytochemical and pharmacological studies of this plant indicate that it produces glycosides, sterols, triterpenes, lactones and flavonoids<sup>1,10</sup>.

In a previous study different chemical constituents were isolated from *B. guianensis*, showing anti-inflammatory and analgesic activities<sup>5,13</sup>. It is known that some compounds of plant origin are produced by their endophytic fungi<sup>12</sup> and that some of the biological activities described for plants can be found in their endophytes<sup>8</sup>.

Thus, the present work is aimed to perform the prospection of the antimicrobial activity of the extracts of endophytic fungi from *B. guianensis* to obtain antimicrobial compounds.

A specimen of *B. guianensis* (voucher specimen No. IAN 177,179) was collected from "Embrapa Amazônia Oriental - Belém, Brazil" and endophytic fungi were isolated after washing with water. A small fragment of plant was subjected to a series of immersions for disinfection and elimination of epiphytic fungi, first in hexane PA (Tedia®) for 1 min, then in 70% ethyl alcohol for 30 s, then in sodium hypochlorite solution 2% for 2 min and finally in sterile water.

After, the plant material was inoculated in Petri dishes containing PDA (Potato, Dextrose, Agar) culture medium for growth of colonies. Seventeen (17) fungi were isolated and purified by successive samplings. Fungi were identified by colony morphology and microscopic aspects observation on optical microscopy and DNA sequencing using the ITS4 region. Small fragments of the isolated fungi were transferred to 10 ml flasks containing distilled water<sup>6</sup> and stored in the mycology collection of the "Laboratório de Biotecnologia e Química de Micro-organismos/Universidade Federal do Pará" (LaBQuiM/UFPA).

Fungi were grown in two 125 ml Erlenmeyer flasks containing 25 g of rice and 10 ml of distilled water each. The flasks were autoclaved at 121 °C for 45 min (Vertical Autoclave 75 L, Prismatec®), and then fungi were inoculated in sterile conditions and incubated at room temperature for 23 days in a static mode for growth of colonies. Then, methanol (Tedia®) was added to the biomass. After 24 h, the material was filtered and the methanolic solutions were evaporated in a rotary evaporator (Quimis®, Q344B, São Paulo, Brazil) to obtain the extracts.

Susceptibility of the microorganisms to the test extracts was determined by the microbroth dilution assay as recommended by the Subcommittee on Antifungal Susceptibility Testing of the United State National Committee for Clinical Laboratory Standards<sup>9</sup>, which was performed on 96 well plates containing 100 µl of Mueller Hinton Broth (MHB), 100 µl of test extracts and 5 µl of test bacteria at  $1.0 \times 10^8$  CFU/ml, followed by incubation at 37 °C (24 h). The test extracts obtained from the fungal culture were dissolved in dimethylsulfoxide at concentration 39–2500 µg/ml. The test microorganisms were *Escherichia coli* (ATCC 25922), *Pseudomonas aeruginosa* (ATCC 27853), *S. aureus* (ATCC 25923), *Bacillus subtilis* (ATCC 6633) and *Salmonella* Typhimurium (ATCC 14028) (provided by the Instituto Evandro Chagas - Belém, Brazil). Bioactivity was recorded as the absence of red color in the wells after the addition of 10 µl of 2,3,5-triphenyltetrazolium chloride. Penicillin, vancomycin and tetracycline (25 µg/ml each) were used as positive controls; the cultivation medium (MHB only) was used as negative control.

None of the extracts showed significant activity against bacteria *P. aeruginosa* and *S. aureus*. With respect to *E. coli*, the extract of the fungus *Exserohilum rostratum* (ER1.1) inhibited growth at a concentration of 78 µg/ml and in the case of bacteria *B. subtilis* the greatest inhibitions were observed for the extracts of fungus *Aspergillus* sp. (EJC08), *Paecilomyces* sp. (EJC01.1) and *E. rostratum* (ER1.1) at a concentration of 78 µg/ml. For *S. Typhimurium* bacteria the best result obtained was for the extracts of fungus *Pestalotiopsis* sp. (EJF02) at a concentration of 156 µg/ml. Based on the results obtained (Table 1) the extract of

**Table 1** Results obtained to antimicrobial assay determined by microbroth dilution with methanolic extracts (2500–39 µg/ml) of endophytic fungi from *B. guianensis*

	Code	Fungus	Minimum inhibitory concentration (µg/ml) bacteria tested				
			<i>Ec</i>	<i>Pa</i>	<i>Bs</i>	<i>Sa</i>	<i>St</i>
1	EJC03	<i>Colletotrichum</i> sp.	312	NA	>2000	NA	NA
2	EJC01.9	<i>C. coccodes</i>	NA	>2000	NA	NA	NA
3	EJF08	<i>Colletotrichum</i> sp.	>2000	NA	NA	312	>2000
4	EJF10	<i>Colletotrichum</i> sp.	NA	NA	NA	SA	NA
5	EJC04	<i>Aspergillus</i> sp.	1250	NA	NA	312 312 <sup>a</sup>	NA
6	EJC08	<i>Aspergillus</i> sp.	NA	>2000	78	NA	>2000
7	EJC07	<i>Pestalotiopsis</i> sp.	>2000	>2000	NA	>2000	>2000
8	EJF02	<i>Pestalotiopsis</i> sp.	650	>2000	>2000	NA	156
9	EJCP12	<i>Scedosporium</i> sp.	>2000	>2000	>2000	1250	>2000
10	EJCP13	<i>S. apiospermum</i>	650	>2000	156	312 1250 <sup>a</sup>	1250
11	EJC10	<i>C. clavata</i>	>2000	NA	1250	NA	>2000
12	EJC11	<i>Phomopsis</i> sp.	156	NA	312	NA	NA
13	ER1.1	<i>E. rostratum</i>	78	312	78	NA	NA
14	EJC01.1	<i>Paecilomyces</i> sp.	156 650 <sup>a</sup>	NA	78 156 <sup>a</sup>	NA	>2000
15	EJCP05	<i>Xylaria</i> sp.	>2000	NA	>2000	NA	NA
16	EJCP07	<i>Xylaria</i> sp.	650	650	650	NA	NA
17	EJCP11	<i>Xylaria</i> sp.	>2000	NA	312	NA	NA

*Ec* = *E. coli*, *Pa* = *P. aeruginosa*, *Bs* = *B. subtilis*, *Sa* = *S. aureus*, *St* = *S. thiphymurium*, NA = no action.


<sup>a</sup> MBC (µg/ml).

fungus *E. rostratum* (ER1.1) was selected for the isolation of the active compound.

The methanolic extract obtained from selected *E. rostratum* fungi (ER 1.1) (2.0g) was fractioned on silica gel column chromatography using hexane/ethyl acetate (9:1, 4:1, 7:3, 1:1, 3:7), ethyl acetate, ethyl acetate/methanol (3:7, 1:1) and methanol resulting in 9 fractions. The fraction hexane/ethyl acetate 7:3 (230 mg) was chromatographed on a silica gel column eluted with hexane/ethyl acetate (9:1, 4:1, 7:3, 1:1, 3:7), ethyl acetate resulting in 85 fractions pooled in F1 to F8; fraction F3 gave a solid orange-colored oil compound 1 (18 mg). Compound 1 was identified by 1D and 2D nuclear magnetic resonance (NMR) and mass spectrometry (MS). After isolation and identification, compound 1 was tested against *E. coli* bacteria (ATCC 25922), *P. aeruginosa* (ATCC 27853), *S. aureus* (ATCC 25923), *B. subtilis* (ATCC 6633) and *S. Typhimurium* (ATCC 14028) at a concentration of 500–7.8 µg/ml.

The <sup>1</sup>H and <sup>13</sup>C NMR experiments were recorded in a NMR spectrometer (Mercury 300, Varian, Oxford, Oxfordshire, UK) with CDCl<sub>3</sub> (Cambridge<sup>®</sup>) as solvent and standard. MS spectra were carried out in the mass spectrometer (Acquity TQD, Waters, Milford, MA, USA) using electrospray ionization in positive ion mode, ESI(+). The specific rotation was performed using specific rotation equipment (Nova Instruments NO 1412, Piracicaba, Brazil).

Compound 1 was isolated as orange oil and was optically active with the specific rotation [α]<sub>D</sub><sup>20</sup>: +60 (c 1.0, dichloromethane). The molecular formula C<sub>16</sub>H<sub>20</sub>O<sub>6</sub> was calculated as base of its mass spectrum ESI(+) *m/z* 309 [M+H]<sup>+</sup>. The <sup>1</sup>H NMR spectrum showed signal to the chelated hydroxyl



**Fig. 1** Compound 1 isolated from the methanol extract of *E. rostratum*.

( $\delta_{\text{H}}$  11.27, OH-8) and signals to an aromatic ring hydrogen ( $\delta_{\text{H}}$  6.59), three carbinolic hydrogens ( $\delta_{\text{H}}$  5.04, 4.53 and 4.11), and two OCH<sub>3</sub> groups ( $\delta_{\text{H}}$  3.94 and 3.88). The analysis of the <sup>13</sup>C NMR and HSQC spectra showed the presence of a carbonyl ester conjugate ( $\delta_{\text{C}}$  167.8) chelated by hydrogen bond with hydroxyl, six carbons of the aromatic ring, three of which were oxygenated ( $\delta_{\text{C}}$  158.7, 156.2 and 137.3) and one unreplaced ( $\delta_{\text{C}}$  104.3), three carbinolic carbons ( $\delta_{\text{C}}$  81.1, 78.7 and 74.4), three methylene ( $\delta_{\text{C}}$  39.0, 38.0 and 19.1), and methyl ( $\delta_{\text{C}}$  13.9). NMR data to compound 1 (Table 2) were consistent with the structure of the isocoumarin derivative, monocerin<sup>15</sup> (Fig. 1).

Antimicrobial testing was performed for compound 1, showing good activity against *E. coli* bacteria (MIC

**Table 2**  $^1\text{H}$  and  $^{13}\text{C}$  NMR data (300 MHz and 75 MHz,  $\text{CDCl}_3$ ) to compound 1

n <sup>o</sup>	H (mult, J in Hz)	C
1		167.8
3	5.04 ( <i>dd</i> , 5.4 and 3.0)	81.1
4	4.53 ( <i>d</i> , 3.0)	74.4
4a		131.1
5	6.59 ( <i>s</i> )	104.3
6		158.7
7		137.3
8		156.2
8a		102.0
9a	2.14 ( <i>ddd</i> , 14.6; 6.0 and 1.1)	39
9b	2.58 ( <i>ddd</i> , 14.6; 8.4 and 6.0)	–
10	4.11 ( <i>m</i> )	78.7
11	1.65 ( <i>m</i> )	38.0
12	1.38 ( <i>m</i> )	19.1
13	0.90 ( <i>t</i> , 7.2)	13.9
OMe-6	3.94 ( <i>s</i> )	56.2
OMe-7	3.88 ( <i>s</i> )	60.7
OH-8	11.27 ( <i>s</i> )	

15.62  $\mu\text{g}/\text{ml}$  and MBC 250  $\mu\text{g}/\text{ml}$ ), *P. aeruginosa* (MIC 15.62  $\mu\text{g}/\text{ml}$  and MBC 62.5  $\mu\text{g}/\text{ml}$ ), *B. subtilis* (MIC 15.62  $\mu\text{g}/\text{ml}$  and MBC 62.5  $\mu\text{g}/\text{ml}$ ), *S. aureus* (MIC 62.5  $\mu\text{g}/\text{ml}$ ) and *S. Typhimurium* (MIC 31.25  $\mu\text{g}/\text{ml}$ ). Data suggest that compound 1 is responsible for the activity observed for the *E. rostratum* extract. Monocerin has been isolated as an antifungal, insecticidal, and phytotoxic secondary metabolite from several fungal species including *Helminthosporium monoceras*, *Exserohilum turcicum*, *Fusarium larvarum* and *Microdochium bolleyi* and showed to be also active against *Plasmodium falciparum*<sup>11,15</sup>. Zhang et al. (2008)<sup>15</sup> isolated monocerin from fungus strain *M. bolleyi* and presented antimicrobial activity against *E. coli* in the agar diffusion test.

The present work isolated 17 endophytic fungi from *B. guianensis*, being the methanolic extract of fungus *E. rostratum* the most active. Monocerin was isolated by chromatographic procedures, showing good antimicrobial activity. This work reports for the first time the isolation of monocerin from endophytic fungi of this Amazon plant, highlighting the activity of this compound as a broad spectrum antimicrobial agent.

## Ethical disclosures

**Protection of human and animal subjects.** The authors declare that the procedures followed were in accordance with the regulations of the relevant clinical research ethics committee and with those of the Code of Ethics of the World Medical Association (Declaration of Helsinki).

**Confidentiality of data.** The authors declare that no patient data appear in this article.

**Right to privacy and informed consent.** The authors declare that no patient data appear in this article.

## Conflict of interest

The authors declare that they have no conflicts of interest.

## Acknowledgments

The authors thank Fundação Amazônia de Amparo a Estudos e Pesquisa do Pará (FAPESPA), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), and Coordenação de Aperfeiçoamento de Pessoal de Ensino Superior (CAPES) for their financial support.

## References

- Achenbach H, Stocker M, Constenla MA. Flavonoid and other constituents of *Bauhinia manca*. *Phytochemistry*. 1988;27:1835–41.
- Aly AH, Debbab A, Kjer J, Proksch P. Fungal endophytes from higher plants: a prolific source of phytochemicals and other bioactive natural products. *Fungal Divers*. 2010;41:1–16.
- Brady SF, Clardy JC. CR377, a new pentaketide antifungal agent isolated from an endophytic fungus. *J Nat Prod*. 2000;63:1447–8.
- Cafêu MC, Silva GH, Teles HL, Araujo AR, Bolzani VS, Young MCM, Pfenning LH. Substâncias antifúngicas de *Xylaria* sp., um fungo endofítico isolado de *Palicourea marcgravii* (rubiaceae). *Quim Nova*. 2005;28:991–5.
- Carvalho JCT, Santos LS, Viana EP, Almeida SSMS, Marconato E, Rodrigues M, Ferreira LR, Van der Kamp A. Anti-inflammatory and analgesic activities of the crude extracts from stem bark of *Bauhinia guianensis*. *Pharm Biol*. 1999;37:281–4.
- Castellani A. Viability of some pathogenic fungi in distilled water. *J Trop Med Hyg*. 1939;42:225–6.
- Dai J, Krohn K, Draeger S, Schulz B. New naphthalene-chroman coupling products from the endophytic fungus *Nodulisporium* sp. from *Erica arborea*. *Eur J Org Chem*. 2009;10:1564–9.
- Li H, Qing C, Zhang Y, Zhao Z. Screening for endophytic fungi with antitumor and antifungal activities from Chinese medicinal plants. *World J Microbiol Biotechnol*. 2005;21:1515–9.
- National Committee for Clinical Laboratory Standards. Methods for dilution and antimicrobial susceptibility tests for bacteria that grow aerobically. Wayne, PA: NCCLS Approved Standard M7-A4; 1997.
- Pinheiro EAA, Carvalho JMC, Santos DCP, Feitosa AO, Marinho PSB, Guilhon GMSP, Santos LS, Souza ALD, Marinho AMR. Chemical constituents of *Aspergillus* sp EJC08 isolated as endophyte from *Bauhinia guianensis* and their antimicrobial activity. *An Acad Bras Cienc*. 2013;85:1247–52.
- Sappapan R, Sommit D, Ngamrojanavanich N, Pengprecha S, Wiyakrutta S, Sriubolmas N, Pudhom K. 11-Hydroxymonocerin from the plant endophytic fungus *Exserohilum rostratum*. *J Nat Prod*. 2008;71:1657–9.
- Strobel GA, Daisy B, Castillo U, Harper J. Natural products from endophytic microorganisms. *J Nat Prod*. 2004;67:257–68.
- Viana EP, Santa-Rosa RS, Almeida SSMS, Santos LS. Constituents of the stem bark of *Bauhinia guianensis*. *Fitoterapia*. 1999;70:111–2.
- World Health Organization. Media centre – antibiotic resistance. <http://www.who.int/mediacentre/factsheets/fs194/en/> [accessed 10.04.15].
- Zhang W, Krohn K, Draeger S, Schulz B. Bioactive isocoumarins isolated from the endophytic fungus *Microdochium bolleyi*. *J Nat Prod*. 2008;71:1078–81.